

AD _____

COOPERATIVE AGREEMENT NO: DAMD17-94-V-4004

TITLE: TESTING OF EXPERIMENTAL ANTILEISHMANIAL COMPOUNDS

PRINCIPAL INVESTIGATOR: William L. Hanson, Ph.D.

CONTRACTING

ORGANIZATION: University of Georgia Research Foundation, Inc.
Boyd Graduate Studies Research Center (GSRC)
The University of Georgia
Athens, Georgia 30602-7411

REPORT DATE: October 19, 1994

TYPE OF REPORT: Final Report

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick
Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19941215 162

THIS DOCUMENT IS UNCLASSIFIED

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 19/10/94	3. REPORT TYPE AND DATES COVERED FINAL REPORT (28/03/94-23/09/94)		
4. TITLE AND SUBTITLE TESTING OF EXPERIMENTAL ANTILEISHMANIAL COMPOUNDS		5. FUNDING NUMBERS Cooperative Agreement No. DAMD17-94-V-4004		
6. AUTHOR(S) Dr. William L. Hanson, Principal Investigator				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The University of Georgia Research Foundation, Inc. Boyd Graduate Studies Research Center (GSRC) The University of Georgia Athens, GA 30602-7411		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U. S. Army Medical Research and Materiel Command Fort Detrick Frederick, MD 21702-5012		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; Distribution unlimited		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) <p>Six plant derivatives which were selected for <u>in vivo</u> study because of their <u>in vitro</u> antileishmanial activity and low toxicity were studied for activity against <u>Leishmania</u> <u>Leishmania donovani</u> and <u>Leishmania Viannia braziliensis</u> in hamsters. The time interval between completion of treatment and evaluation of results was extended up to 4-6 weeks to evaluate any possible delayed antileishmanial activity by the compounds. None of the compounds studied were active in hamsters against either species of <u>Leishmania</u>. Extension of time between treatment and evaluation of results did not increase the <u>in vivo</u> activity of any of the compounds studied.</p> <p>None of the compounds studied were toxic based on body weight loss, mortality, or clinical signs.</p>				
14. SUBJECT TERMS <u>Leishmania</u> <u>Leishmania donovani</u> ; <u>Leishmania Viannia braziliensis</u> ; Chemotherapy; Plant derivatives		15. NUMBER OF PAGES 14		
		16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

I
FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

_____ Where copyrighted material is quoted, permission has been obtained to use such material.

_____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

_____ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

X In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

_____ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

_____ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

_____ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

X In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

William L. Hanson Oct. 19 1994
PI - Signature Date

ACKNOWLEDGEMENTS

I would like to thank Department of Parasitology, College of Veterinary Medicine personnel, Ms. Nell H. Landers, Office Manager, for administrative and clerical assistance and Ms. Barbara L. Harris, Laboratory Technician II, for technical assistance with this study. Their efforts are appreciated.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

TABLE OF CONTENTS

Foreword.....	1
Acknowledgements.....	2
Introduction.....	4
Materials and Methods.....	5
I. Visceral Test System.....	5
II. Cutaneous Test System.....	6
Results.....	7
Discussion.....	7
Conclusions.....	7
Literature Cited.....	8
Appendix.....	9
Table I.....	10
Table II.....	11
Personnel Employed from this Cooperative Agreement.....	12
Bibliography of Published Work.....	12
Graduate Degrees Awarded.....	12
Distribution List.....	13

INTRODUCTION

Protozoan parasites of the genus *Leishmania* are widespread throughout the world where they cause a complex of visceral or cutaneous diseases in human beings as well as some animals including dogs in numerous tropical and sub-tropical countries (1,2,3). Since the leishmaniasis commonly exist as zoonoses, these diseases pose a significant potential threat to military personnel as well as military dogs throughout endemic areas. Relatively recent publicity regarding infection of personnel involved in Operation Desert Storm has reemphasized the military significance of the leishmaniasis.

Better drugs are needed for the treatment of the leishmaniasis since those currently available are often not satisfactorily effective and are potentially toxic to man and animals.

This laboratory has been involved for several years in studies to identify new compounds for antileishmanial activity against both visceral (*Leishmania donovani*) and cutaneous (*Leishmania Viannia braziliensis*) leishmaniasis. Although several new compounds have been identified with activity against *L. (V.) braziliensis*, none have shown adequate promise to warrant initiation of clinical trials. However, among the most promising active compounds found against visceral leishmaniasis during these studies is the 8-aminoquinoline, WR06026 (4). This compound is now undergoing clinical trials in Kenyan visceral leishmaniasis patients. Testing for compounds active against visceral leishmaniasis has continued during this project period in the event that WR06026 does not perform in the field as expected and testing has continued to identify more active and less toxic compounds for *L. (V.) braziliensis*. Emphasis was placed on the study of *in vivo* activity of plant derivatives which have been noted previously to be active *in vitro* against *Leishmania*.

This report summarizes the results of studies conducted for this contract during the period March 28, 1994 through September 23, 1994.

MATERIALS AND METHODS

I. Visceral Test System

A Khartoum strain of *L. (L.) donovani* (WR378) was used and the golden hamster (*Mesocricetus auratus*), 50-70 gm, served as the host animal. Suspensions of amastigotes for infection of experimental hamsters were prepared by grinding heavily infected hamster spleens in sterile saline in a Ten Broeck tissue grinder and diluting the suspensions so that 0.2 ml contained approximately 10×10^6 amastigotes. Each experimental hamster was infected via the intracardiac injection of 0.2 ml of the amastigote suspension.

The testing procedure used was that described by Stauber and his associates (5,6,7) as modified by Hanson, et al. (8) with the exception that the time interval between completion of treatment and termination of the experiment was extended (see next paragraph). On Day 3 following infection, hamsters were divided randomly into experimental groups consisting of a minimum of 6 animals per group, initial group weights were obtained, and administration of test compounds was initiated. Each compound was tested at 2 or 3 drug dosage levels dependent on the priority rating and nature of the compound.

The vehicle for the reference and test compounds was 0.6% DMSO. Each test group contained six hamsters. A control group of six hamsters received the 0.6% DMSO vehicle only and the reference compound, Glucantime®, was given at 2 drug dosage levels, 416 and 208 mg sb/kg. All test compounds were administered at 832 total mg/kg. The reference compound, Glucantime®, was administered via the intramuscular route and the test compounds were administered orally twice daily on days 3 through 6. Final group weights were obtained on all experimental hamsters on Day 7 and all hamsters were killed four weeks after completion of treatment, livers removed, weighed, and liver impressions made for enumeration of amastigotes. Subsequently, the total number of parasites per liver was determined as described by Stauber, et al. (5,6,7).

In addition to recording body weight changes as a general indicator of toxicity of the test compounds, experimental hamsters were observed for such clinical signs of toxicity as nervous disorders, roughened hair coat, and sluggish activity. Deaths of the animals was also considered indicative of significant drug toxicity.

After determining the ratio of numbers of amastigotes per host cell nucleus, the weight of the organ, and initial and final weights of the hamsters, the raw data was evaluated with a Gateway 2000 microcomputer using a program which calculates percent weight change, total numbers of parasites, mean numbers of parasites per organ, and percent parasite suppression. The computer program then performs linear and non-linear regression analysis and calculates an SD_{50} for active compound from each of the analyses (drug dosage resulting in 50% suppression of amastigotes). The SD_{50} from the non-linear analysis is used for a comparison of the relative efficacy of the test compounds and the efficacy of test compounds relative to that of the reference compound, Glucantime®. The linear regression analysis is included only for comparison with the non-linear analysis.

II. Cutaneous Test System

Leishmania (V.) braziliensis (WR539) was used in these studies. Male golden hamsters, 50-70 gm, served as experimental hosts.

Promastigotes for establishing experimental infections in hamsters were grown in Schneider's *Drosophila* Medium (Hendricks, et al., 9) and quantitated using procedures described previously (Hanson and Roberson, 10). In preparation for infection and weekly during the experiment, the hair was clipped on the dorsal tail head and a commercial depilatory agent applied to the area to remove the remaining hair. Each hamster was inoculated via the intradermal route with approximately 1.5×10^7 promastigotes of *L. (V.) braziliensis* near the base of the tail using a 0.25 ml glass syringe equipped with a 30 gauge X 1/2" needle. Each experimental group consisted of six hamsters. Initial body weights were obtained and administration of therapy was initiated on Day 19 postinfection, and continued through Day 22 postinfection. Glucantime® was included at two dosage levels (832 and 208 total mg/Sb/kg) as the reference compound, and a group of six hamsters received vehicle only (0.6% DMSO). Test compounds were administered at 780 total mg/kg. The reference compound, Glucantime®, was administered via the intramuscular route and the test compounds were administered orally.

Lesion area of each experimental hamster was determined one week after completion of treatment and again at eight weeks after completion of treatment with the aid of a template made at WRAIR and calibrated according to the formula $r_1 r_2 \pi$ where r_1 is the major radius of the lesion and r_2 is the minor radius (Wilson, et al., 11). The mean lesion area of each experimental group was obtained and the percent suppression of lesion size calculated by comparing the mean lesion area of each treated group with that of the group receiving vehicle only with the aid of a computer program and a Gateway 2000 microcomputer. The computer program performs linear and non-linear regression analysis and calculates an SD_{50} for each active compound using both analyses. The SD_{50} obtained from the non-linear analyses is used for a rough comparison of the relative efficacies of the test compounds and the relative efficacy of test compounds with that of the reference compound, Glucantime®. The linear regression analysis is performed for comparison with the non-linear analysis.

RESULTS

Six plant derivatives which have been noted by others to be active *in vitro* against *Leishmania* were noted to be inactive in hamsters when tested against either *L. (L.) donovani* (Table I) or *L. (V.) braziliensis* (Table II). Extension of time up to 4-6 weeks between treatment and evaluation of results did not increase the *in vivo* activity of any of the compounds studied in either test system.

None of the compounds were toxic to hamsters as indicated by mortality, weight loss, or clinical signs.

DISCUSSION

The purposes of these studies were (1) to determine whether selected plant derivatives with *in vitro* antileishmanial activity were also active *in vivo* in hamsters, and (2) to determine if extending the interval between completion of therapy and evaluation of parasite numbers or lesion size would reveal possible evidence of delayed activity of the test compounds.

Although these compounds had been observed to be active *in vitro*, no evidence of *in vivo* antileishmanial activity was noted in our studies. It is not unusual for test compounds to be active against parasites *in vivo* and not be active *in vitro* which may be due to a variety of factors (12).

Extension of the time interval between administration of the test compounds and evaluation of their effect did not reveal any evidence of delayed activity of the test compounds.

CONCLUSIONS

1. The probability of identifying compounds with *in vivo* antileishmanial activity is low.
2. Although the compounds selected for these studies were active *in vitro* but not *in vivo*, *in vitro* activity must remain a major criterion for selecting test compounds.
3. Although the plant derivatives selected for these studies were not active, this approach has tremendous merit and should continue since historically considerable success has been achieved with plant derivatives against such parasites as amoebae, *Plasmodium*, and various helminths.

LITERATURE CITED

1. Kinnamon, K. E., E. A. Steck, P. S. Loizeaux, L. D. Hendricks, V. B. Waits, W. L. Chapman, Jr., and W. L. Hanson. 1979. Leishmaniasis: Military significance and new hope for treatment. Mil. Med. 44(10): 660-664.
2. Tropical Diseases Research, Seventh Programme Report, 1 January 1983 - 31 December 1984. UNDP/World Bank/WHO Imprimerie A. Barthelemy, Avignon, France 1985. pp 7/3-7/18.
3. Chapman, Willie L., Jr. and William L. Hanson. 1984. Leishmaniasis. IN Clinical Microbiology and Infectious Diseases of the Dog and Cat. W. B. Saunders Company, Philadelphia. pp 764-770.
4. Kinnamon, Kenneth E., Edgar A. Steck, Peter S. Loizeaux, William L. Hanson, Willie L. Chapman, Jr. and Virginia B. Waits. 1978. The antileishmanial activity of lepidines. Am. J. Trop. Med. Hyg. 27(4): 751-757.
5. Stauber, L. A., E. M. Franchino, and J. Grun. 1958. An eight-day method for screening compounds against *Leishmania donovani* in the golden hamster. J. Protozool. 5: 269-273.
6. Stauber, L. A. 1958. Host resistance to the Khartoum strain of *Leishmania donovani*. The Rice Institute Pamphlet Vol. XLV(1): 80-96.
7. Stauber, L. A. 1958. Chemotherapy of experimental leishmaniasis. Proc. 6th International Congr. on Trop. Med. and Mal. III: 797-805.
8. Hanson, W. L., W. L. Chapman, Jr., and K. E. Kinnamon. 1977. Testing of drugs for antileishmanial activity in golden hamsters infected with *Leishmania donovani*. Internat'l. J. Parasitol. 7: 443-447.
9. Hendricks, L. D., D. Wood, and M. Hajduk. 1978. Hemoflagellates: Commercially available liquid media for rapid cultivation. Parasitol. 76: 309-316.
10. Hanson, W. L. and E. L. Roberson. 1974. Density of parasites in various organs and the relation to number of trypomastigotes in the blood during acute infections of *Trypanosoma cruzi* in mice. J. Protozool. 21: 512-517.
11. Wilson, H. R., B. W. Dieckmann, and G. E. Childs. 1979. *Leishmania braziliensis* and *Leishmania mexicana*: Experimental cutaneous infections in golden hamsters. Exptl. Parasitol. 47: 270-283.
12. Hudson, A. T. 1994. The continuation of empiricism to antiparasite drug discovery. Parasitology Today 10(10): 387-389.

Appendix

Table I. Summary of the suppressive activity of selected compounds against *Leishmania Leishmania donovani* four weeks after the completion of treatment in the golden hamster.

Treatment	Total mg/kg	Route	Percent Suppression
Vehicle (DMSO)	-	PO	-
BL09186 (Glucantime®)	416 208	IM	36 21
BN38070	832	PO	2
BN38089	832	PO	-23*
BN38098	832	PO	-5*
BN38105	832	PO	1
BN38114	832	PO	2
BN38123	832	PO	-7*

* Negative percent suppression signifies a greater mean number of amastigotes/liver in the experimental group than in the vehicle control group.

PO: per os
IM: intramuscular

Table II. Summary of the suppressive activity of selected compounds against *Leishmania Viannia braziliensis* one week and eight weeks after completion of treatment in the golden hamster.

Treatment	Total mg/kg	Route	Percent Suppression	
			1 week	8 weeks
Vehicle (DMSO)	-	PO	-	-
BL09186 (Glucantime®)	832 208	IM	62 39	26 16
BN38070	780	PO	26	3
BN38089	780	PO	16	3
BN38098	780	PO	10	6
BN38105	780	PO	-10*	-6*
BN38114	780	PO	10	-12*
BN38123	780	PO	-13*	-10*

* Negative percent suppression signifies a larger mean lesion area in the experimental group than in the vehicle control group.

PO: per os
IM: intramuscular

Personnel Employed from this Cooperative Agreement

<u>Name and Position</u>	<u>Percent Effort</u>	<u>Length of Employment</u>
Virginia B. Waits Research Coordinator II	50%	07/01/94 - 07/31/94
Virginia B. Waits Research Coordinator II	57%	08/01/94 - 08/31/94
Virginia B. Waits Research Coordinator II	100%	09/01/94 - 09/30/94
Virginia B. Waits Research Coordinator II	.476 of 1% (1 Full Day 8.0 hours)	10/01/94 - 10/01/94

Bibliography of Published Work

None

Graduate Degrees Resulting from this Cooperative Agreement

None

Distribution List

Six (6) Copies: U. S. Army Medical Research and Materiel Command
(Provisional), ATTN: SGRD-RMI-S/Virginia Miller
Fort Detrick
Frederick, Maryland 21702-5012

One (1) Copy: Ms. Linda B. Allen
Director
Sponsored Programs
The University of Georgia Research Foundation, Inc.
The University of Georgia
Athens, GA 30602-7411